

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

1,3 -disubstituted urea derivatives, useful as inhibitors of acyl  
co-enzyme A cholesterol acyl transferase enzyme

Patent Assignee: SLOVAKOFARMA AS (SLOV-N)

Inventor: FABEROVA V; KAKALIK I; OREMUS V; SCHMIDTOVA L; SMAHOVSKY V;  
ZEMANEK M

Number of Countries: 083 Number of Patents: 005

Patent Family:

Patent No Kind Date Applicat No Kind Date Week

WO 9932437 A1 19990701 WO 98SK19 A 19981216 199933 B

SK 9701751 A3 19990712 SK 971751 A 19971219 199939

AU 9916976 A 19990712 AU 9916976 A 19981216 199950

EP 1042278 A1 20001011 EP 98961715 A 19981216 200052

WO 98SK19 A 19981216

CZ 200002297 A3 20001115 WO 98SK19 A 19981216 200064

CZ 20002297 A 19981216

Priority Applications (No Type Date): SK 971751 A 19971219

Patent Details:

Patent No Kind Lan Pg Main IPC Filing Notes

WO 9932437 A1 E 33 C07C-275/36

Designated States (National): AL AM AT AU AZ BA BB BG BR BY CA CH CN CU

CZ DE DK EE ES FI GB GE GH GM HR HU ID IL IN IS JP KE KG KP KR KZ LC LK

LR LS LT LU LV MD MG MK MN MW MX NO NZ PL PT RO RU SD SE SG SI SK SL TJ

TM TR TT UA UG US UZ VN YU ZW

Designated States (Regional): AT BE CH CY DE DK EA ES FI FR GB GH GM GR

IE IT KE LS LU MC MW NL OA PT SE UG ZW

SK 9701751 A3 C07C-273/18

AU 9916976 A C07C-275/36 Based on patent WO 9932437

EP 1042278 A1 E C07C-275/36 Based on patent WO 9932437

Designated States (Regional): AT BE CH DE DK ES FI FR GB GR IE IT LI LU

NL PT RO SE SI

CZ 200002297 A3 C07C-275/30 Based on patent WO 9932437

Abstract (Basic): WO 9932437 A1

NOVELTY - Novel 1,3-disubstituted ureas are inhibitors of acyl  
co-enzyme A: cholesterol acyl transferase (ACAT) enzyme, and can  
inhibit cholesterol esterification and absorption in  
hypercholesterolemia.

DETAILED DESCRIPTION - 1,3-disubstituted ureas of formula (I) are  
new.

When R1=2-fluorophenyl, 2,4- or 2,5-difluorophenyl, 2-chlorophenyl,  
2,3-, 2,6- or 3,5-dichlorophenyl, 2- or 4-methylphenyl, 2,4-, 2,6- or  
3,5-dimethylphenyl, 2,6-di(methylethyl)phenyl, 2-, 3- or  
4-trifluoromethylphenyl, 2-, 3- or 4-pyridyl, 1- or 2-naphthyl or

1-adamantyl, then R2=NO2, and X=O;

when R1=groups as above and 4-nitrophenyl, 4-fluorophenyl, 4-chlorophenyl, 2,4- or 3,4-dichlorophenyl, then R2=NO2 and X=S;

when R1=2,4-difluorophenyl, 2,3-dichlorophenyl, or 2,6-di(methylethyl)phenyl, then R2=amino; and X=O or S.

ACTIVITY - Anti-hypercholesterolemia.

MECHANISM OF ACTION - ACAT inhibitors. The percentages of ACAT inhibition by compounds (I), at concentration 2 mM, was up to 88% in rats' liver, and up to 71% in rabbits' intestinal mucosa.

USE - (I) are used for treating hypercholesterolemia and preventing atherosclerosis.

pp; 33 DwgNo 0/0

Title Terms: DISUBSTITUTED; UREA; DERIVATIVE; USEFUL; INHIBIT; ACYL; CO; ENZYME; CHOLESTEROL; ACYL; TRANSFERASE; ENZYME

Derwent Class: B05

International Patent Class (Main): C07C-273/18; C07C-275/30; C07C-275/36

International Patent Class (Additional): A61K-031/17; A61K-031/44;

A61P-009/00; C07C-275/00; C07C-275/28; C07C-275/34; C07C-323/44;

C07D-213/75; A61K-031-44



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification <sup>6</sup> : C07C 275/36, 323/44, C07D 213/75 // A61K 31/17, 31/44</p>	<p>A1</p>	<p>(11) International Publication Number: WO 99/32437 (43) International Publication Date: 1 July 1999 (01.07.99)</p>
<p>(21) International Application Number: PCT/SK98/00019 (22) International Filing Date: 16 December 1998 (16.12.98) (30) Priority Data: PV 1751-97 19 December 1997 (19.12.97) SK (71) Applicant (for all designated States except US): SLOVAKO-FARMA, A.S. [SK/SK]; Železničná 12, 920 27 Hlohovec (SK). (72) Inventors; and (75) Inventors/Applicants (for US only): OREMUS, Vladimír [SK/SK]; Vysoká 4, 811 06 Bratislava (SK). SMAHOVSKÝ, Vendelín [SK/SK]; Novomeského 18, 902 01 Pezinok (SK). FÁBEROVÁ, Viera [SK/SK]; Račianska 12, 831 04 Bratislava (SK). KAKALÍK, Ivan [SK/SK]; 900 81 Šenkvice (SK). SCHMIDTOVÁ, Ľudmila [SK/SK]; Komenského 10, 900 01 Modra (SK). ZEMÁNEK, Marián [SK/SK]; Zochova 16, 811 03 Bratislava (SK). (74) Agent: NEUSCHL, Jozef; Rott, Ružička and Guttmann, Patentová, známková a právna kancelária, v.o.s., Pionierska 15, 831 02 Bratislava (SK).</p>		<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published With international search report. With amended claims.</p>
<p>(54) Title: 1,3-DISUBSTITUTED UREAS AS ACAT INHIBITORS, AND METHOD OF PREPARING THEREOF</p> <p>(57) Abstract</p> <p>The invention relates to 1,3-disubstituted ureas of general formula (I) where R<sup>1</sup> is an aryl, R<sup>2</sup> is nitro and/or amino, and X is oxygen and/or sulfur, and the method of preparing thereof which consists in treating aromatic amines with isocyanates. Isocyanates may be formed in situ and the reaction carried out in toluene, at 80 °C. If the nitro group is formed, it is reduced with hydrogen in the presence of palladium catalyst to the amino group. The obtained 1,3-disubstituted ureas are inhibitors of the activity of the acyl co-enzyme A: cholesterol acyltransferase (ACAT) enzyme, and may be used to inhibit cholesterol esterification and absorption in hypercholesterolemia.</p> <div style="text-align: center;"> <p>(I)</p> </div>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LJ	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

**1,3-Disubstituted Ureas as ACAT Inhibitors, and Method of Preparing Thereof****Technical Field**

The invention relates to compounds the principal characteristics of which include inhibition of the acyl-coenzyme A: cholesterol acyltransferase (ACAT) enzyme activity, and to a method for the preparation of such compounds.

**Background Art**


The acyl-coenzyme A: cholesterol O-acyltransferase (EC 2.3.1.26) (ACAT) enzyme is responsible for the catalysis of the intracellular esterification of cholesterol. ACAT is present in most tissues such as the intestine, liver, and arterial wall. The enzyme is assumed to be involved in numerous processes which underlie the development of atherosclerosis, absorption of dietary cholesterol, accumulation of cholesterol esters, hepatic secretion of cholesterol esters into the blood plasma in the form of VLDL cholesterol.

A number of substances of the urea type have been described to inhibit ACAT. We shall show several more recent examples describing 1,3-disubstituted ureas as ACAT enzyme inhibitors. Patents EP 506532, FR 2674522, JP 93097802, US 5219859 describe ureas containing indole derivatives in their molecules. A combination of aromatic and aliphatic moieties has been described in Patents EP 665216, JP 95258199. Introduction into the molecule of a 1,3-dioxolane ring has been reported in Bioorg. Med. Chem. Lett. 1995, 5(15): 1581.

1,3-Disubstituted ureas of the present invention have not been described in literature.

**Disclosure of Invention**

1,3-Disubstituted ureas of general formula I,



wherein R<sup>1</sup> is 4-nitrophenyl, 2-fluorophenyl, 4-fluorophenyl, 2,4-difluorophenyl, 2,5-difluorophenyl, 2,6-difluorophenyl, 2-chlorophenyl, 4-chlorophenyl, 2,3-dichlorophenyl, 2,4-dichlorophenyl, 2,6-dichlorophenyl, 3,4-dichlorophenyl, 3,5-dichlorophenyl, 2-methylphenyl, 4-methylphenyl, 2,4-dimethylphenyl, 2,6-dimethylphenyl, 3,5-dimethylphenyl, 2,6-di(methylethyl)phenyl, 2-trifluoromethylphenyl, 3-trifluoromethylphenyl, 4-trifluoromethylphenyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 1-naphthyl, 2-naphthyl, 1-adamantyl, and R<sup>2</sup> is nitro, and X = O, S; and for R<sup>1</sup> being 2,4-difluorophenyl, 2,3-dichlorophenyl, 2,6-dimethylphenyl, 2,6-di(methylethyl)-phenyl R<sup>2</sup> is amino, and X = O, S.

The method for the preparation of the above compounds according to this invention consists in reacting an isocyanate (as prepared in situ or as commercially available) with amine to give an urea the nitro group of which may subsequently be reduced to the amino group. Ureas prepared in this way show inhibitory effect on acyl-coenzyme A: cholesterol acyltransferase (ACAT).

**Examples****Example 1**

1-(4-Nitrophenyl)-3-((-4 nitrophenoxy)-phenyl)-urea

A solution of 4-nitrophenylisocyanate in diethylether (20 ml) is added dropwise to a solution of 4'-nitrophenoxy-aniline (2.30 g, 0.01 mol) in a mixture of diethylether (20 ml) and tetrahydrofurane (20 ml) at laboratory temperature, and the mixture is stirred for 16 hours. The precipitated product is aspirated, washed with diethylether (20 ml). The raw product is purified by chromatography on silica gel eluting with dichloromethane-methanol.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): 7.11(d, 2H, H-arom.); 7.17(d, 2H, H-arom.); 7.59(d, 2H, H-arom.); 7.70(d, 2H, H-arom.); 8.20(d, 2H, H-arom.); 7.25(d, 2H, H-arom.); 9.05(s, 1H, NH.); 9.46(s, 1H, NH).

<sup>13</sup>C-NMR (CDCl<sub>3</sub>): 116.80(CH-arom.); 117.46(CH-arom.); 120.46(CH-arom.); 121.13(CH-arom.); 125.07(CH-arom.); 126.11(CH-arom.); 136.50, 141.00, 141.96, 146.26, 148.86, 151.95(C-arom.); 163.39(C=O).

Analysis for C<sub>18</sub>H<sub>14</sub>N<sub>4</sub>O<sub>6</sub>

%C(calcd/found)	%H	%N
57.85/57.73	3.58/3.61	14.21/14.12

Yield: 92% Melting temp.: 231-234°C

#### Example 2

##### 1-(2-Fluorophenyl)-3-((4'-nitrophenoxy)-phenyl)-urea

The title compound was prepared from 2-fluorophenylisocyanate by an analogous procedure to that described in Example 1.

Analysis for C<sub>18</sub>H<sub>14</sub>FN<sub>3</sub>O<sub>4</sub>

%C(calcd/found)	%H	%N
62.11/62.09	3.84/3.88	11.44/11.29

Yield: 48% Melting temp.: 253-255°C

#### Example 3

##### 1-(4-Fluorophenyl)-3-((4'-nitrophenoxy)-phenyl)-urea

The title compound was prepared from 4-fluorophenylisocyanate by an analogous procedure to that described in Example 1.

Analysis for  $C_{18}H_{14}FN_3O_4$

%C(calcd/found)	%H	%N
62.11/61.99	3.84/3.85	11.44/11.34

Yield: 59% Melting temp.: 267-269°C

#### Example 4

1-(2,4-Difluorophenyl)-3-((4'-nitrophenoxy)-phenyl)-urea

The title compound was prepared from 2,4-difluorophenylisocyanate by an analogous procedure to that described in Example 1.

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ ): 6.98-7.18(m, 5H, H-arom.); 7.23-7.37(m, 1H, H-arom.); 7.56(d, 2H, H-arom.); 8.03-8.16(m, 1H, H-arom.); 8.24(d, 2H, H-arom.); 8.50(s, 1H, NH); 9.13(s, 1H, NH).

$^{13}\text{C-NMR}$  ( $\text{CDCl}_3$ ): 103.72(CH-arom.); 110.96(CH-arom.); 116.73(CH-arom.); 119.88(CH-arom.); 121.18(CH-arom.); 122.02(CH-arom.); 126.09(CH-arom.); 136.94, 141.91, 148.48, 152.27, 154.49, 159.80(C-arom.); 163.46(C=O).

Analysis for  $C_{18}H_{13}F_2N_3O_4$

%C(calcd/found)	%H	%N
59.22/59.20	3.40/3.53	10.91/10.89

Yield: 85% Melting temp.: 223-224°C

#### Example 5

1-(2,5-Difluorophenyl)-3-((4'-nitrophenoxy)-phenyl)-urea

The title compound was prepared from 2,5-difluorophenylisocyanate by an analogous procedure to that described in Example 1.

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ ): 6.73-6.88(m, 1H, H-arom.); 7.04-7.35(m, 5H, H-arom.); 7.56(d, 2H, H-arom.); 7.98-8.11(m, 1H, H-arom.); 8.22(d, 2H, H-arom.); 8.75-9.30(br.s., 2H, NH).

$^{13}\text{C-NMR}$  ( $\text{CDCl}_3$ ): 106.56(CH-arom.); 107.75(CH-arom.); 115.67(CH-arom.); 116.74(CH-arom.); 120.03(CH-arom.); 121.21(CH-arom.); 126.08(CH-arom.); 128.82, 136.62, 141.95, 148.73, 151.95, 155.65, 160.38(C-arom.); 163.42(C=O).

Analysis for  $C_{19}H_{13}F_2N_3O_4$

%C(calcd/found)	%H	%N
59.22/59.07	3.40/3.49	10.91/10.83

Yield: 76% Melting temp.: 207-208°C

#### Example 6

1-(2,6-Difluorophenyl)-3-((4'-nitrophenoxy)-phenyl)-urea

The title compound was prepared from 2,6-difluorophenylisocyanate by an analogous procedure to that described in Example 1.

$^1H$ -NMR ( $CDCl_3$ ): 6.95-7.26(m, 6H, H-arom.); 7.42-7.56(m, 2H, H-arom.); 8.01-8.23(m, 3H, H-arom.); 8.94-9.05(m, 2H, NH).

$^{13}C$ -NMR ( $CDCl_3$ ): 111.77(CH-arom.); 116.82(CH-arom.); 120.06(CH-arom.); 121.18(CH-arom.); 126.19(CH-arom.); 127.11(CH-arom.); 137.31, 141.99, 148.49, 152.61, 155.65, 160.57(C-arom.); 163.61(C=O).

Analysis for  $C_{19}H_{13}F_2N_3O_4$

%C(calcd/found)	%H	%N
59.22/59.10	3.40/3.55	10.91/10.78

Yield: 75% Melting temp.: 231-232°C

#### Example 7

1-(2-Chlorophenyl)-3-((4'-nitrophenoxy)-phenyl)-urea

The title compound was prepared from 2-chlorophenylisocyanate by an analogous procedure to that described in Example 1.

Analysis for  $C_{19}H_{14}ClN_3O_4$

%C(calcd/found)	%H	%N	%Cl
59.46/59.37	3.68/3.82	10.95/10.78	9.24/8.99

Yield: 63% Melting temp.: 195-197°C

#### Example 8

1-(4-Chlorophenyl)-3-((4'-nitrophenoxy)-phenyl)-urea

The title compound was prepared from 4-chlorophenylisocyanate by an analogous procedure to that described in Example 1.

Analysis for  $C_{19}H_{14}ClN_3O_4$

%C(calcd/found)	%H	%N	%Cl
59.46/59.37	3.68/3.77	10.95/10.84	9.24/9.02

Yield: 69% Melting temp.: 234-236°C

#### Example 9.

1-(2,3-Dichlorophenyl)-3-((4'-nitrophenoxy)-phenyl)-urea

The title compound was prepared from 2,3-dichlorophenylisocyanate by an analogous procedure to that described in Example 1.

Analysis for  $C_{19}H_{13}Cl_2N_3O_4$

%C(calcd/found)	%H	%N	%Cl
54.56/54.50	3.13/3.31	10.05/9.78	16.95/16.91

Yield: 74% Melting temp.: 199-201°C

#### Example 10

1-(2,4-Dichlorophenyl)-3-((4'-nitrophenoxy)-phenyl)-urea

The title compound was prepared from 2,4-dichlorophenylisocyanate by an analogous procedure to that described in Example 1.

Analysis for  $C_{19}H_{13}Cl_2N_3O_4$

%C(calcd/found)	%H	%N	%Cl
54.56/54.49	3.13/3.21	10.05/9.80	16.95/16.59

Yield: 71% Melting temp.: 267-269°C

#### Example 11

1-(2,6-Dichlorophenyl)-3-((4'-nitrophenoxy)-phenyl)-urea

The title compound was prepared from 2,6-dichlorophenylisocyanate by an analogous procedure to that described in Example 1.

Analysis for  $C_{19}H_{13}Cl_2N_3O_4$

%C(calcd/found)	%H	%N	%Cl
54.56/54.39	3.13/3.20	10.05/9.92	16.95/16.81

Yield: 68% Melting temp.: 195-198°C

## Example 12

1-(3,4-Dichlorophenyl)-3-((4'-nitrophenoxy)-phenyl)-urea

The title compound was prepared from 3,4-dichlorophenylisocyanate by an analogous procedure to that described in Example 1.

Analysis for  $C_{18}H_{13}Cl_2N_3O_4$

%C(calcd/found)	%H	%N	%Cl
54.56/54.49	3.13/3.23	10.05/9.89	16.95/16.78

Yield: 80% Melting temp.: 179-180°C

## Example 13

1-(3,5-Dichlorophenyl)-3-((4'-nitrophenoxy)-phenyl)-urea

The title compound was prepared from 3,5-dichlorophenylisocyanate by an analogous procedure to that described in Example 1.

Analysis for  $C_{18}H_{13}Cl_2N_3O_4$

%C(calcd/found)	%H	%N	%Cl
54.56/54.48	3.13/3.30	10.05/10.01	16.95/17.24

Yield: 56% Melting temp.: 213-216°C

## Example 14

1-(2-Methylphenyl)-3-((4'-nitrophenoxy)-phenyl)-urea

The title compound was prepared from 2-methylphenylisocyanate by an analogous procedure to that described in Example 1.

Analysis for  $C_{20}H_{17}N_3O_4$

%C(calcd/found)	%H	%N
60.11/65.96	4.72/4.89	11.56/11.48

Yield: 59% Melting temp.: 112-116°C

## Example 15

1-(4-Methylphenyl)-3-((4'-nitrophenoxy)-phenyl)-urea

The title compound was prepared from 4-methylphenylisocyanate by an analogous procedure to that described in Example 1.

Analysis for  $C_{20}H_{17}N_3O_4$

%C(calcd/found)	%H	%N
66.11/66.02	4.72/4.87	11.56/11.40

Yield: 64% Melting temp.: 168-170°C

#### Example 16

##### 1-(2,4-Dimethylphenyl)-3-((4'-nitrophenoxy)-phenyl)-urea

The title compound was prepared from 2,4-dimethylphenylisocyanate by an analogous procedure to that described in Example 1.

Analysis for  $C_{21}H_{19}N_3O_4$

%C(calcd/found)	%H	%N
66.83/66.87	5.07/5.10	11.13/11.05

Yield: 73% Melting temp.: 165-169°C

#### Example 17

##### 1-(2,6-Dimethylphenyl)-3-((4'-nitrophenoxy)-phenyl)-urea

The title compound was prepared from 2,6-dimethylphenylisocyanate by an analogous procedure to that described in Example 1.

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ ): 2.22(s, 6H,  $\text{CH}_3$ ); 7.04-7.17(m, 7H, H-arom.); 7.57(m, 2H, H-arom.); 7.74(s, 1H, NH); 8.23(d, 2H, H-arom.); 8.87(s, 1H, NH).

$^{13}\text{C-NMR}$  ( $\text{CDCl}_3$ ): 18.19(2x  $\text{CH}_3$ ); 116.61(CH-arom.); 119.54(CH-arom.); 121.08(CH-arom.); 126.07(CH-arom.); 127.67(CH-arom.); 135.53(CH-arom.); 125.93, 135.51, 137.95, 141.82, 147.87, 153.10(CH-arom.); 163.64(C=O).

Analysis for  $C_{21}H_{19}N_3O_4$

%C(calcd/found)	%H	%N
66.83/66.67	5.07/5.18	11.13/10.98

Yield: 68% Melting temp.: 249-250°C

#### Example 18

##### 1-(3,5-Dimethylphenyl)-3-((4'-nitrophenoxy)-phenyl)-urea

The title compound was prepared from 3,5-dimethylphenylisocyanate by an analogous procedure to that described in Example 1.

Analysis for  $C_{21}H_{19}N_3O_4$

%C(calcd/found)	%H	%N
66.83/66.78	5.07/5.22	11.13/11.06

Yield: 58% Melting temp.: 145-147°C

#### Example 19

1-(2,6-Di-(methylethyl)-phenyl)-3-((4'-nitrophenoxy)-phenyl)-urea

The title compound was prepared from 2,6-di-(methylethyl)phenylisocyanate by an analogous procedure to that described in Example 1.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): 1.18(d, 6H, 2xCH<sub>3</sub>); 3.22(hept., 2H, 2xCH); 7.06-7.30(m, 7H, H-arom.); 7.56(d, 2H, H-arom.); 7.63(s, 1H, NH); 8.23(d, 2H, H-arom.); 8.78(s, 1H, NH).

<sup>13</sup>C-NMR (CDCl<sub>3</sub>): 23.31(2xCH<sub>3</sub>); 27.86(2xCH); 116.58(CH-arom.); 119.44(CH-arom.); 120.86(CH-arom.); 122.72(CH-arom.); 125.87(CH-arom.); 146.53(CH-arom.); 127.08, 132.14, 137.80, 141.83, 147.89, 154.10(C-arom.); 163.46(C=O).

Analysis for C<sub>26</sub>H<sub>27</sub>N<sub>3</sub>O<sub>4</sub>

%C(calcd/found)	%H	%N
69.27/69.13	6.28/6.34	9.69/9.56

Yield: 88% Melting temp.: 208-210°C

#### Example 20

1-(2-Trifluoromethylphenyl)-3-((4'-nitrophenoxy)-phenyl)-urea

4'-Nitrophenoxy-aniline (1.0 g, 4.3 mmol), triphosgene (0.43 g, 1.44 mmol), triethylamine (0.6 ml, 4.3 mmol) are heated in toluene (15 ml) in a pressure tube at 80°C for 20 hours. Then, 2-trifluoromethylaniline (0.53 ml, 4.3 mmol) and triethylamine (0.6 ml, 4.3 mmol) in toluene (10 ml) are added. The mixture is heated at 80°C for 4 hours, then it is concentrated, and the product is isolated using chromatography on silica gel eluting with dichloromethane - methanol.

Analysis for C<sub>20</sub>H<sub>14</sub>F<sub>3</sub>N<sub>3</sub>O<sub>4</sub>

%C(calcd/found)	%H	%N
57.56/57.60	3.38/3.44	10.07/9.89

Yield: 67% Melting temp.: 201-203°C

**Example 21****1-(3-Trifluoromethylphenyl)-3-((4'-nitrophenoxy)-phenyl)-urea**

The title compound was prepared from 3-trifluoromethylaniline by an analogous procedure to that described in Example 20.

Analysis for  $C_{20}H_{14}F_3N_3O_4$

%C(calcd/found)	%H	%N
57.56/57.66	3.38/3.45	10.07/9.96

Yield: 72% Melting temp.: 208-211°C

**Example 22****1-(4-Trifluoromethylphenyl)-3-((4'-nitrophenoxy)-phenyl)-urea**

The title compound was prepared from 4-trifluoromethylaniline by an analogous procedure to that described in Example 20.

Analysis for  $C_{20}H_{14}F_3N_3O_4$

%C(calcd/found)	%H	%N
57.56/57.48	3.38/3.41	10.07/10.00

Yield: 45% Melting temp.: 185-189°C

**Example 23****1-(2-Pyridyl)-3-(4'-nitrophenoxy)-phenyl)-urea**

The title compound was prepared from 2-pyridylamine by an analogous procedure to that described in Example 20.

Analysis for  $C_{18}H_{14}N_4O_4$

%C(calcd/found)	%H	%N
61.71/61.67	4.03/4.06	15.99/15.79

Yield: 76% Melting temp.: 143-146°C

**Example 24****1-(3-Pyridyl)-3-(4'-nitrophenoxy)-phenyl)-urea**

The title compound was prepared from 3-pyridylamine by an analogous procedure to that described in Example 20.

Analysis for  $C_{18}H_{14}N_4O_4$

%C(calcd/found)	%H	%N
61.71/61.58	4.03/4.21	15.99/15.87

Yield: 69% Melting temp.: 177-179°C

#### Example 25

##### 1-(4-Pyridyl)-3-(4'-nitrophenoxy)-phenyl)-urea

The title compound was prepared from 4-pyridylamine by an analogous procedure to that described in Example 20.

Analysis for  $C_{18}H_{14}N_4O_4$

%C(calcd/found)	%H	%N
61.71/61.66	4.03/4.11	15.99/15.87

Yield: 72% Melting temp.: 126-127°C

#### Example 26

##### 1-(1-Naphthyl)-3-(4'-nitrophenoxy)-phenyl)-urea

The title compound was prepared from 1-naphthylisocyanate by an analogous procedure to that described in Example 1.

Analysis for  $C_{23}H_{17}N_3O_4$

%C(calcd/found)	%H	%N
69.17/69.23	4.29/4.41	10.52/10.46

Yield: 82% Melting temp.: 117-119°C

#### Example 27

##### 1-(2-Naphthyl)-3-(4'-nitrophenoxy)-phenyl)-urea

The title compound was prepared from 1-naphthylamine by an analogous procedure to that described in Example 20.

Analysis for  $C_{23}H_{17}N_3O_4$

%C(calcd/found)	%H	%N
69.17/69.09	4.29/4.36	10.52/10.38

Yield: 69% Melting temp.: 103-106°C

## Example 28

## 1-(1-Adamantyl)-3-(4'-nitrophenoxy)-phenyl)-urea

The title compound was prepared from 1-adamantylamine by an analogous procedure to that described in Example 20.

Analysis for  $C_{23}H_{25}N_3O_4$ .

%C(calcd/found)	%H	%N
67.80/67.65	6.18/6.23	10.31/10.16

Yield: 61% Melting temp.: 143-146°C

## Example 29

## 1-(4-Nitrophenyl)-3-((4'-nitrophenylthio)-phenyl)-urea

A solution of 4-nitrophenylisocyanate in diethylether (20 ml) is added dropwise to a solution of 4'-nitrophenylthio-aniline (2.46 g, 0.01 mol) in a mixture of diethylether (20 ml) and tetrahydrofurane (20 ml) at laboratory temperature, and the mixture is stirred for 16 hours. The resulting product is aspirated, washed with diethylether (20 ml). The crude product is purified by chromatography on silica gel eluting with dichloromethane and methanol.

Analysis for  $C_{19}H_{14}N_4O_5S$

%C(calcd/found)	%H	%N	S%
55.61/55.52	3.44/3.49	13.65/13.59	7.81/7.67

Yield: 56% Melting temp.: 164-167°C

## Example 30

## 1-(2-Fluorophenyl)-3-((4'-nitrophenylthio)-phenyl)-urea

The title compound was prepared from 2-fluorophenylisocyanate by an analogous procedure to that described in Example 29.

Analysis for  $C_{19}H_{14}FN_3O_3S$

%C(calcd/found)	%H	%N	S%
59.52/59.41	3.68/3.77	10.96/11.04	8.36/8.41

Yield: 61% Melting temp.: 274-277°C

**Example 31**

1-(4-Fluorophenyl)-3-((4'-nitrophenylthio)-phenyl)-urea

The title compound was prepared from 4-fluorophenylisocyanate by an analogous procedure to that described in Example 29.

Analysis for  $C_{19}H_{14}FN_3O_3S$

%C(calcd/found)	%H	%N	S%
59.52/59.46	3.68/3.71	10.96/10.87	8.36/8.18

Yield: 59% Melting temp.: 287-290°C

**Example 32**

1-(2,4-Difluorophenyl)-3-((4'-nitrophenylthio)-phenyl)-urea

The title compound was prepared from 2,4-difluorophenylisocyanate by an analogous procedure to that described in Example 29.

Analysis for  $C_{19}H_{13}F_2N_3O_3S$

%C(calcd/found)	%H	%N	S%
56.86/56.78	3.26/3.39	10.47/10.41	7.99/7.86

Yield: 57% Melting temp.: 268-271°C

**Example 33**

1-(2,5-Difluorophenyl)-3-((4'-nitrophenylthio)-phenyl)-urea

The title compound was prepared from 2,5-difluorophenylisocyanate by an analogous procedure to that described in Example 29.

Analysis for  $C_{19}H_{13}F_2N_3O_3S$

%C(calcd/found)	%H	%N	S%
56.86/56.76	3.26/3.37	10.47/10.35	7.99/8.05

Yield: 64% Melting temp.: 259-261°C

**Example 34**

1-(2,6-Difluorophenyl)-3-((4'-nitrophenylthio)-phenyl)-urea

The title compound was prepared from 2,6-difluorophenylisocyanate by an analogous procedure to that described in Example 29.

Analysis for  $C_{19}H_{13}F_2N_3O_3S$

%C(calcd/found)	%H	%N	S%
56.86/56.69	3.26/3.34	10.47/10.43	7.99/7.81

Yield: 68% Melting temp.: 263-265°C

#### Example 35

1-(2-Chlorophenyl)-3-((4'-nitrophenylthio)-phenyl)-urea

The title compound was prepared from 2-chlorophenylisocyanate by an analogous procedure to that described in Example 29.

Analysis for  $C_{18}H_{14}ClN_3O_3S$

%C(calcd/found)	%H	%N	%Cl	S%
57.07/57.01	3.53/3.62	10.51/11.46	8.87/8.65	8.02/7.95

Yield: 65% Melting temp.: 231-233°C

#### Example 36

1-(4-Chlorophenyl)-3-((4'-nitrophenylthio)-phenyl)-urea

The title compound was prepared from 4-chlorophenylisocyanate by an analogous procedure to that described in Example 29.

Analysis for  $C_{19}H_{14}ClN_3O_3S$

%C(calcd/found)	%H	%N	%Cl	S%
57.07/56.97	3.53/3.57	10.51/10.45	8.87/8.81	8.02/7.86

Yield: 63% Melting temp.: 206-209°C

#### Example 37

1-(2,3-Dichlorophenyl)-3-((4'-nitrophenylthio)-phenyl)-urea

The title compound was prepared from 2,3-dichlorophenylisocyanate by an analogous procedure to that described in Example 29.

Analysis for  $C_{19}H_{13}Cl_2N_3O_3S$

%C(calcd/found)	%H	%N	%Cl	S%
52.55/52.46	3.02/3.07	9.68/9.62	16.33/16.27	7.38/7.50

Yield: 75% Melting temp.: 157-159°C

**Example 38**

1-(2,4-Dichlorophenyl)-3-((4'-nitrophenylthio)-phenyl)-urea

The title compound was prepared from 2,4-dichlorophenylisocyanate by an analogous procedure to that described in Example 29.

Analysis for  $C_{19}H_{13}Cl_2N_3O_3S$

%C(calcd/found)	%H	%N	%Cl	S%
52.55/52.57	3.02/3.21	9.68/9.54	16.33/16.35	7.38/7.28

Yield: 57% Melting temp.: 174-178°C

**Example 39**

1-(2,6-Dichlorophenyl)-3-((4'-nitrophenylthio)-phenyl)-urea

The title compound was prepared from 2,6-dichlorophenylisocyanate by an analogous procedure to that described in Example 29.

Analysis for  $C_{18}H_{13}Cl_2N_3O_3S$

%C(calcd/found)	%H	%N	%Cl	S%
52.55/52.51	3.02/3.07	9.68/9.73	16.33/16.25	7.38/7.19

Yield: 83% Melting temp.: 164-167°C

**Example 40**

1-(3,4-Dichlorophenyl)-3-((4'-nitrophenylthio)-phenyl)-urea

The title compound was prepared from 3,4-dichlorophenylisocyanate by an analogous procedure to that described in Example 29.

Analysis for  $C_{19}H_{13}Cl_2N_3O_3S$

%C(calcd/found)	%H	%N	%Cl	S%
52.55/52.47	3.02/3.14	9.68/9.57	16.33/16.09	7.38/7.24

Yield: 57% Melting temp.: 238-240°C

**Example 41**

1-(3,5-Dichlorophenyl)-3-((4'-nitrophenylthio)-phenyl)-urea

The title compound was prepared from 3,5-dichlorophenylisocyanate by an analogous procedure to that described in Example 29.

Analysis for  $C_{19}H_{13}Cl_2N_3O_3S$

%C(calcd/found)	%H	%N	%Cl	S%
52.55/52.47	3.02/3.11	9.68/9.59	16.33/16.21	7.38/7.41

Yield: 67% Melting temp.: 185 - 188°C

#### Example 42

##### 1-(2-Methylphenyl)-3-((4'-nitrophenylthio)-phenyl)-urea

The title compound was prepared from 2-methylphenylisocyanate by an analogous procedure to that described in Example 29.

Analysis for  $C_{20}H_{17}N_3O_3S$

%C(calcd/found)	%H	%N	S%
63.31/63.22	4.52/4.66	11.07/10.79	8.45/8.34

Yield: 78% Melting temp.: 229-234°C

#### Example 43

##### 1-(4-Methylphenyl)-3-((4'-nitrophenylthio)-phenyl)-urea

The title compound was prepared from 4-methylphenylisocyanate by an analogous procedure to that described in Example 29.

Analysis for  $C_{20}H_{17}N_3O_3S$

%C(calcd/found)	%H	%N	S%
63.31/63.25	4.52/4.63	11.07/11.12	8.45/8.35

Yield: 73% Melting temp.: 163-166°C

#### Example 44

##### 1-(2,4-Dimethylphenyl)-3-((4'-nitrophenylthio)-phenyl)-urea

The title compound was prepared from 2,4-dimethylphenylisocyanate by an analogous procedure to that described in Example 29.

Analysis for  $C_{21}H_{19}N_3O_3S$

%C(calcd/found)	%H	%N	S%
64.11/64.08	4.87/4.83	10.68/10.59	8.15/7.95

Yield: 65% Melting temp.: 209-213°C

**Example 45**

1-(2,6-Dimethylphenyl)-3-((4'-nitrophenylthio)-phenyl)-urea

The title compound was prepared from 2,6-dimethylphenylisocyanate by an analogous procedure to that described in Example 29.

Analysis for  $C_{21}H_{19}N_3O_3S$

%C(calcd/found)	%H	%N	S%
64.11/63.97	4.87/4.83	10.68/10.47	8.15/8.01

Yield: 72% Melting temp.: 264-267°C

**Example 46**

1-(3,5-Dimethylphenyl)-3-((4'-nitrophenylthio)-phenyl)-urea

The title compound was prepared from 3,5-dimethylphenylisocyanate by an analogous procedure to that described in Example 29.

Analysis for  $C_{21}H_{18}N_3O_3S$

%C(calcd/found)	%H	%N	S%
64.11/64.04	4.87/4.99	10.68/10.63	8.15/8.01

Yield: 68% Melting temp.: 194-196°C

**Example 47**

1-(2,6-Dimethylethyl)-phenyl-((4'-nitrophenylthio)-phenyl)-urea

The title compound was prepared from 2,6-(methylethyl)-phenylisocyanate by an analogous procedure to that described in Example 29.

Analysis for  $C_{25}H_{27}N_3O_3S$

%C(calcd/found)	%H	%N	S%
66.79/66.72	6.06/6.17	9.35/9.27	7.12/6.96

Yield: 72% Melting temp.: 175-177°C

**Example 48**

1-(2-Trifluoromethylphenyl)-3-((4'-nitrophenylthio)-phenyl)-urea

4'-Nitrophenylthio-aniline (1.06 g, 4.3 mmol), triphosgene (0.43 g, 1.44 mmol), triethylamine (0.6 g, 4.3 mmol) in toluene (15 ml) are heated in a pressure tube at 80°C for 20 hours. Subsequently, 2-trifluoromethylaniline (0.53 ml, 4.3 mmol)

and triethylamine (0.6 ml, 4.3 mmol) in toluene (10 ml) are added. The mixture is heated at 80°C for 4 hours, then concentrated, and the product is separated by chromatography on silica gel eluting with dichloromethane-methanol.

Analysis for  $C_{20}H_{14}F_3N_3O_3S$

%C(calcd/found)	%H	%N	S%
55.43/55.38	3.26/3.39	9.70/9.71	7.40/7.35

Yield: 52% Melting temp.: 257-261°C

#### Example 49

1-(3-Trifluoromethylphenyl)-3-((4'-nitrophenylthio)-phenyl)-urea

The title compound was prepared from 3-trifluorophenylisocyanate by an analogous procedure to that described in Example 48.

Analysis for  $C_{20}H_{14}F_3N_3O_3S$

%C(calcd/found)	%H	%N	S%
55.43/55.21	3.26/3.38	9.70/9.53	7.40/7.31

Yield: 65% Melting temp.: 241-244°C

#### Example 50

1-(4-Trifluoromethylphenyl)-3-((4'-nitrophenylthio)-phenyl)-urea

The title compound was prepared from 4-trifluorophenylisocyanate by an analogous procedure to that described in Example 48.

Analysis for  $C_{20}H_{14}F_3N_3O_3S$

%C(calcd/found)	%H	%N	S%
55.43/55.37	3.26/3.33	9.70/9.81	7.40/7.28

Yield: 51% Melting temp.: 254-257°C

#### Example 51

1-(2-Pyridyl)-3-((4'-nitrophenylthio)-phenyl)-urea

The title compound was prepared from 2-pyridylamine by an analogous procedure to that described in Example 48.

Analysis for  $C_{18}H_{14}N_4O_3S$

%C(calcd/found)	%H	%N	S%
59.01/58.86	3.85/3.91	15.29/15.23	8.75/8.80

Yield: 48% Melting temp.: 278-281°C

#### Example 52

1-(3-Pyridyl)-3-((4'-nitrophenylthio)-phenyl)-urea

The title compound was prepared from 3-pyridylamine by an analogous procedure to that described in Example 48.

Analysis for  $C_{18}H_{14}N_4O_3S$

%C(calcd/found)	%H	%N	S%
59.01/58.99	3.85/3.99	15.29/15.25	8.75/8.49

Yield: 63% Melting temp.: 261-264°C

#### Example 53

1-(4-Pyridyl)-3-((4'-nitrophenylthio)-phenyl)-urea

The title compound was prepared from 4-pyridylamine by an analogous procedure to that described in Example 48.

Analysis for  $C_{18}H_{14}N_4O_3S$

%C(calcd/found)	%H	%N	S%
59.01/58.92	3.85/3.76	15.29/15.32	8.75/8.67

Yield: 69% Melting temp.: 190-192°C

#### Example 54

1-(1-Naphthyl)-3-((4'-nitrophenylthio)-phenyl)-urea

The title compound was prepared from 1-naphthylisocyanate by an analogous procedure to that described in Example 29.

Analysis for  $C_{23}H_{17}N_3O_3S$

%C(calcd/found)	%H	%N	S%
66.49/66.53	4.13/4.21	10.12/10.17	7.70/7.54

Yield: 56% Melting temp.: 164-168°C

## Example 55

## 1-(2-Naphthyl)-3-((4'-nitrophenylthio)-phenyl)-urea

The title compound was prepared from 2-naphthylamine by an analogous procedure to that described in Example 48.

Analysis for  $C_{23}H_{17}N_3O_3S$

%C(calcd/found)	%H	%N	S%
66.49/66.47	4.13/4.25	10.12/10.06	7.70/7.57

Yield: 69% Melting temp.: 142-147°C

## Example 56

## 1-(1-Adamantyl)-3-((4'-nitrophenylthio)-phenyl)-urea

The title compound was prepared from 1-adamantylamine by an analogous procedure to that described in Example 48.

Analysis for  $C_{23}H_{25}N_3O_3S$

%C(calcd/found)	%H	%N	S%
65.22/65.17	5.95/6.03	9.93/10.02	7.56/7.38

Yield: 52% Melting temp.: 264-267°C

## Example 57

## 1-(2,4-Difluorophenyl)-3-((4'-aminophenoxy)-phenyl)-urea

One gram of compound 4 is dissolved in methanol (20 ml) and 0.1 g of 10% palladium on charcoal is added. The mixture is stirred under hydrogen atmosphere (at atmospheric pressure) for 20 hours. Subsequently, 100 ml methanol is added and the catalyst is removed by filtering. The product is then obtained by concentrating the methanolic solution.

Analysis for  $C_{19}H_{15}F_2N_3O_2$

%C(calcd/found)	%H	%N
64.22/64.09	4.25/4.29	11.83/12.01

Yield: 88% Melting temp.: 248-251°C

## Example 58

## 1-(2,5-Dichlorophenyl)-3-((4'-aminophenoxy)-phenyl)-urea

The title compound was prepared from compound 9 by an analogous procedure to that described in Example 57.

Analysis for  $C_{18}H_{15}Cl_2N_3O_2$

%C(calcd/found)	%H	%N	%Cl
58.78/58.45	3.89/3.94	10.82/10.78	18.26/18.11

Yield: 91% Melting temp.: 201-204°C

#### Example 59

1-(2,6-Dimethylphenyl)-3-((4'-aminophenoxy)-phenyl)-urea

The title compound was prepared from compound 17 by an analogous procedure to that described in Example 57.

Analysis for  $C_{21}H_{21}N_3O_2$

%C(calcd/found)	%H	%N
72.60/72.45	6.09/6.13	12.10/12.02

Yield: 85% Melting temp.: 225-227°C

#### Example 60

1-(2,6-Di(methylethyl)-phenyl)-3-((4'-aminophenoxy)-phenyl)-urea

The title compound was prepared from compound 19 by an analogous procedure to that described in Example 57.

$^1H$ -NMR ( $CDCl_3$ ): 1.15(d, 6H, 2xCH<sub>3</sub>); 3.15(hept., 2H, 2xCH); 4.86(s, 2H, NH); 6.57(d, 2H, H-arom.); 6.72(d, 2H, H-arom.); 6.81(d, 2H, H-arom.); 7.10-1.28(m, 3H, H-arom.); 7.36(d, 2H, H-arom.); 7.56(s, 1H, HN); 8.58(br.s., 1H, NH).

$^{13}C$ -NMR ( $CDCl_3$ ): 23.43(2xCH<sub>3</sub>); 27.93(2xCH); 114.63(CH-arom.); 117.55(CH-arom.); 119.09(CH-arom.); 199.43(CH-arom.); 122.79(CH-arom.); 127.11(CH-arom.); 127.11(CH-arom.); 132.40, 134.86, 144.77, 146.61, 146.89, 152.88(C-arom.); 154.36(C=O).

Analysis for  $C_{26}H_{29}N_3O_2$

%C(calcd/found)	%H	%N
74.41/74.54	7.24/7.33	10.41/10.34

Yield: 90% Melting temp.: 219-221°C

**Example 61**

1-(2,4-Difluorophenyl)-3-((4'-aminophenylthio)-phenyl)-urea

The title compound was prepared from compound 32 by an analogous procedure to that described in Example 57.

Analysis for  $C_{19}H_{15}F_2N_3OS$

%C(calcd/found)	%H	%N	%S
61.44/61.56	4.07/4.18	11.31/11.15	8.63/8.51

Yield: 79% Melting temp.: exceeding 300°C

**Example 62**

1-(2,3-Dichlorophenyl)-3-((4'-aminophenylthio)-phenyl)-urea

The title compound was prepared from compound 37 by an analogous procedure to that described in Example 57.

Analysis for  $C_{19}H_{15}Cl_2N_3OS$

%C(calcd/found)	%H	%N	%Cl	%S
56.44/56.35	3.74/3.80	10.39/10.41	17.54/17.57	7.93/7.59

Yield: 88% Melting temp.: 259-261°C

**Example 63**

1-(2,6-Dimethylphenyl)-3-((4'-aminophenylthio)-phenyl)-urea

The title compound was prepared from compound 45 by an analogous procedure to that described in Example 57.

Analysis for  $C_{21}H_{21}N_3OS$

%C(calcd/found)	%H	%N	%S
69.39/69.32	5.82/5.93	11.56/11.49	8.28/8.54

Yield: 94% Melting temp.: 198-202°C

**Example 64**

1-(2,6-Di-(methylethyl)-phenyl)-3-((4'-aminophenylthio)-phenyl)-urea

The title compound was prepared from compound 47 by an analogous procedure to that described in Example 57.

Analysis for  $C_{25}H_{28}N_3OS$ 

%C(calcd/found)	%H	%N	%S
71.56/71.55	6.97/6.89	10.01/10.11	7.64/7.58

Yield: 83% Melting temp.: 267-271°C

**Tests**

The biological activity of the substances was evaluated based on the in vitro inhibition of acylCoA:cholesterol acyltransferase (ACAT) activity. The enzyme was obtained from the microsomal fraction of rat liver cells and rabbit intestinal mucosa of animals fed with cholesterol. The substrates for the enzyme reaction included exogenous oleoyl co-enzyme A and endogenous cholesterol.  $^{14}C$ -oleoyl co-enzyme A conversion to  $^{14}C$ -cholesteryl oleate was monitored. From the mixture of extracted lipids, cholesteryl oleate was separated using thin-layer chromatography, and was quantified radiometrically. ACAT specific activity was expressed as the amount of cholesteryl oleate formed per minute per mg microsomal protein.

Table 1 shows percentages of ACAT inhibition in the rat liver and the rabbit intestinal mucosa at various concentrations of the substances tested. Efficiency was calculated as compared to enzyme activity measured in the presence of 1% dimethylsulfoxide used as the solvent to prepare solutions of the substances tested.

Table 1

Inhibitory effect on rat liver and rabbit intestinal mucosa ACAT activity

No	Efficiency	(%)	Concentration	No	Efficiency	(%)	Concentration
	<i>liver</i>	<i>mucosa</i>	( $\mu$ M)		<i>liver</i>	<i>mucosa</i>	( $\mu$ M)
1	0	46	2	33	0	16	2
2	15	32	2	34	11	25	2
3	0	24	2	35	20	26	2
4	37	55	2	36	12	21	2
5	49	58	2	37	0	0	2
6	0	42	2	38	12	16	2
7	0	0	2	39	15	21	2
8	17	13	2	40	0	20	2
9	58	51	2	41	0	0	2
10	10	26	2	42	27	31	2
11	20	25	2	43	21	32	2
12	11	57	2	44	19	31	2
13	0	0	2	45	23	34	2
14	0	0	2	46	18	27	2
15	0	0	2	47	88	71	2
16	0	0	2	48	25	38	2
17	41	65	2	49	34	45	2
18	0	0	2	50	23	25	2
19	50	67	2	51	48	46	2
20	46	42	2	52	45	36	2
21	38	45	2	53	53	35	2
22	25	18	2	54	24	36	2
23	26	34	2	55	16	31	2
24	0	0	2	56	45	56	2
25	11	17	2	57	53	64	2
26	14	22	2	58	55	46	2


27	0	12	2	59	38	62	2
28	43	58	2	60	68	64	2
29	0	34	2	61	22	25	2
30	0	23	2	62	15	26	2
31	25	27	2	63	21	29	2
32	16	23	2	64	56	67	2

### **Industrial Applicability**

The compounds according to the invention and the method of preparing thereof can be used in pharmaceutical production to make preparations with inhibitory effect on the enzyme acyl co-enzyme A and on cholesterol absorption in hypercholesterolemia.


## CLAIMS

1. 1,3-Disubstituted ureas of general formula I,



wherein  $R^1$  is 4-nitrophenyl, 2-fluorophenyl, 4-fluorophenyl, 2,4-difluorophenyl, 2,5-difluorophenyl, 2,6-difluorophenyl, 2-chlorophenyl, 4-chlorophenyl, 2,3-dichlorophenyl, 2,4-dichlorophenyl, 2,6-dichlorophenyl, 3,4-dichlorophenyl, 3,5-dichlorophenyl, 2-methylphenyl, 4-methylphenyl, 2,4-dimethylphenyl, 2,6-dimethylphenyl, 3,5-dimethylphenyl, 2,6-di(methylethyl)phenyl, 2-trifluoromethylphenyl, 3-trifluoromethylphenyl, 4-trifluoromethylphenyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 1-naphthyl, 2-naphthyl, 1-adamantyl, and  $R^2$  is nitro, and  $X = O, S$ ; and for  $R^1$  being 2,4-difluorophenyl, 2,3-dichlorophenyl, 2,6-dimethylphenyl, 2,6-di(methylethyl)-phenyl  $R^2$  is amino, and  $X = O, S$ .

2. A method of preparing 1,3-disubstituted ureas of general formula I according to claim 1, characterized in that an amine of general formula II,



wherein  $R^2$  and X have the above defined meanings, is treated with an isocyanate of general formula III,



III

wherein  $R^1$  has the above defined meaning, said isocyanate optionally being formed in situ from appropriate reactants,

thus giving the above defined urea.

3. The method of claim 2, characterized in that when said isocyanate is formed in situ, the reaction is carried out in toluene at about 80°C.


4. The method of any of the preceding claims, characterized in that the obtained 1,3-disubstituted urea of general formula I wherein  $R^2$  means nitro, is treated with hydrogen in the presence of palladium catalyst to reduce the nitro group to the amino group.

5. 1,3-Disubstituted ureas of general formula I, according to claim 1 and/or prepared by the method of claim 2 to 4, characterized in that they have inhibitory effect on the acyl co-enzyme A:cholesterol acyltransferase (ACAT) enzyme.

## AMENDED CLAIMS

[received by the International Bureau on 08 June 1999 (08.06.99);  
original claim 1 amended remaining claims unchanged (2 pages)]

1. 1,3-Disubstituted ureas of general formula I,




wherein R<sup>1</sup> is 2-fluorophenyl, 2,4-difluorophenyl, 2,5-difluorophenyl, 2,6-difluorophenyl, 2-chlorophenyl, 2,3-dichlorophenyl, 2,6-dichlorophenyl, 3,5-dichlorophenyl, 2-methylphenyl, 4-methylphenyl, 2,4-dimethylphenyl, 2,6-dimethylphenyl, 3,5-dimethylphenyl, 2,6-di(methylethyl)phenyl, 2-trifluoromethylphenyl, 3-trifluoromethylphenyl, 4-trifluoromethylphenyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 1-naphthyl, 2-naphthyl, 1-adamantyl, and R<sup>2</sup> is nitro, and X = O;

wherein R<sup>1</sup> is 4-nitrophenyl, 2-fluorophenyl, 4-fluorophenyl, 2,4-difluorophenyl, 2,5-difluorophenyl, 2,6-difluorophenyl, 2-chlorophenyl, 4-chlorophenyl, 2,3-dichlorophenyl, 2,4-dichlorophenyl, 2,6-dichlorophenyl, 3,4-dichlorophenyl, 3,5-dichlorophenyl, 2-methylphenyl, 4-methylphenyl, 2,4-dimethylphenyl, 2,6-dimethylphenyl, 3,5-dimethylphenyl, 2,6-di(methylethyl)phenyl, 2-trifluoromethylphenyl, 3-trifluoromethylphenyl, 4-trifluoromethylphenyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 1-naphthyl, 2-naphthyl, 1-adamantyl, and R<sup>2</sup> is nitro, and X = S;

and for R<sup>1</sup> being 2,4-difluorophenyl, 2,3-dichlorophenyl, 2,6-dimethylphenyl, 2,6-di(methylethyl)-phenyl R<sup>2</sup> is amino, and X = O, S.


2. A method of preparing 1,3-disubstituted ureas of general formula I according to claim 1, characterized in that an amine of general formula II,

AMENDED SHEET (ARTICLE 19)



II

wherein  $R^2$  and X have the above defined meanings, is treated with an isocyanate of general formula III,



III

wherein  $R^1$  has the above defined meaning, said isocyanate optionally being formed in situ from appropriate reactants,

thus giving the above defined urea.

3. The method of claim 2, characterized in that when said isocyanate is formed in situ, the reaction is carried out in toluene at about 80°C.

4. The method of any of the preceding claims, characterized in that the obtained 1,3-disubstituted urea of general formula I wherein  $R^2$  means nitro, is treated with hydrogen in the presence of palladium catalyst to reduce the nitro group to the amino group.

5. 1,3-Disubstituted ureas of general formula I, according to claim 1 and/or prepared by the method of claim 2 to 4, characterized in that they have inhibitory effect on the acyl co-enzyme A:cholesterol acyltransferase (ACAT) enzyme.

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/SK 98/00019

**A. CLASSIFICATION OF SUBJECT MATTER**  
IPC 6 C07C275/36 C07C323/44 C07D213/75 //A61K31/17,A61K31/44

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)  
IPC 6 C07C C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 3 284 433 A (H. J. BECKER ET AL) 8 November 1966 see claims 6,8 see examples	1,2
X	EP 0 709 225 A (NIPPON PAPER INDUSTRIES CO) 1 May 1996 see page 15, line 35	1

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

31 March 1999

Date of mailing of the international search report

12/04/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Van Geyt, J

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/SK 98/00019

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 3284433	A	08-11-1966	CH 459174 A	
			DE 1468337 A	28-11-1968
			FR 1469459 A	10-05-1967
			GB 1068887 A	
<hr/>				
EP 0709225	A	01-05-1996	JP 8118806 A	14-05-1996
			JP 8156407 A	18-06-1996
			CA 2161376 A	28-04-1996
			DE 69503864 D	10-09-1998
			DE 69503864 T	18-03-1999
			US 5710094 A	20-01-1998
<hr/>				